

# Asthma First Aid

## 1 Sit the person upright

- Be calm and reassuring
- Do not leave them alone



## 2 Give 4 separate puffs of blue/grey reliever puffer

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer

Repeat until 4 puffs have been taken

**Remember: Shake, 1 puff, 4 breaths**

OR give 2 separate doses of a Bricanyl inhaler (age 6 & over) or a Symbicort inhaler (over 12)



## 3 Wait 4 minutes

- If there is no improvement, give 4 more separate puffs of blue/grey reliever as above

OR give 1 more dose of Bricanyl or Symbicort inhaler



## 4 If there is still no improvement call emergency assistance Dial Triple Zero

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every 4 minutes until emergency assistance arrives

OR give 1 dose of a Bricanyl or Symbicort every 4 minutes - up to 3 more doses of Symbicort



### Call emergency assistance immediately - Dial Triple Zero (000)

- If the person is not breathing
- If the person's asthma suddenly becomes worse or is not improving
- If the person is having an asthma attack and a reliever is not available
- If you are not sure if it's asthma
- If the person is known to have Anaphylaxis - follow their Anaphylaxis Action Plan, then give Asthma First Aid

**Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma**



## Asthma Australia

Contact your local Asthma Foundation  
**1800 ASTHMA Helpline** (1800 278 462)  
[asthmaaustralia.org.au](http://asthmaaustralia.org.au)

© Asthma Australia 2016 Supported by the Australian Government



Translating and  
Interpreting Service  
131 450